

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ


BEST AVAILABLE COPY

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.




Signed

*[Handwritten signature]*

Dated

2 March 2004

Patents Act 1977  
(Rule 16)



# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

24 MAR 2003

1. Your reference JIM/LJ/P20717GB

2. Patent application number  
(The Patent Office will fill in this part) 0306736.0

3. Full name, address and postcode of the or of each applicant (underline all surnames)

NESTEC SA  
CASE POSTALE 353  
CH-1800 VEVEY  
SWITZERLAND

4255063002

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

SWITZERLAND

4. Title of the invention

GELLED FOOD WITH A LIQUID CENTRE

5. Name of your agent (if you have one) ELKINGTON AND FIFE

"Address for service" in the United Kingdom to which all correspondence should be sent  
(including the postcode)

ELKINGTON AND FIFE  
PROSPECT HOUSE  
8 PEMBROKE ROAD  
SEVENOAKS  
KENT  
TN13 1XR

Patents ADP number (if you know it)

67004 ✓

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of Filing (day/month/year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of Filing (day/month/year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer "Yes" if:

- a) any applicant named in part 3 is not an inventor, or  
b) there is an inventor who is not named as an applicant, or  
c) any named applicant is a corporate body.

YES

See note (d))

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form -

Description 14

Claim(s) 4

Abstract 1

Drawing(s) 2 + 2

10. If you are also filing any of the following, state how many against each item.

Priority documents -

Translations of priority documents -

Statement of inventorship and right to grant of a patent (Patents Form 7/77) -

Request for preliminary examination and search (Patents Form 9/77) 1

Request for substantive examination (Patents Form 10/77) -

Any other documents (please specify) -

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

24 March 2003

12. Name and daytime telephone number of person to contact in the United Kingdom

Mr J I MARCHANT  
01732 458881

## GELLED FOOD PRODUCT WITH A LIQUID CENTRE

### FIELD OF THE INVENTION

- 5 This invention is related to a food product comprising a gelled water-based shell and a liquid centre.

### BACKGROUND OF THE INVENTION

- 10 Gums and jellies constituted by water-based hydrocolloid gels represent a significant and growing proportion of the sugar based confectionery market (see Sugar Confectionery Manufacture edited by E B Jackson, Blackie 1990, pages 190 - 217 Chapter 10, Gums and Jellies by E T Best). Such products are usually solid and examples include pastilles, gummy bears, jelly babies and the like. However, liquid
- 15 filled gelled products are also known and examples include Rowntrees Fruit Rush, Rowntrees Bursting Bugs and Betty Crocker Fruit Gushers. The shells of these products, which are often referred to as "liquid-filled gummies" are aqueous based hydrocolloid gels and are believed to be based on starch and/or gelatin.
- 20 The use of gelatin in food products has significant disadvantages. Food-grade gelatin is obtained from bovine or porcine raw materials and is not acceptable to certain groups on religious grounds or to vegetarians. Concerns have also been expressed as to the use of bovine products as a result of bovine spongiform encephalopathy ("BSE").
- 25 A hydrocolloid which has been suggested as a replacement for gelatin in the formation of gelled products is carrageenan. Carrageenans are generally obtained by extraction from seaweeds and are polysaccharides consisting of alternating copolymers of  $\beta(1\rightarrow3)$ -D-galactose and  $\alpha(1\rightarrow4)$ -3,6-anhydro-D-galactose units. At least 8 types of carrageenan are known differing in the amount of sulphate ester and/or other
- 30 substituent groups (see Stortz & Cerezo, Carbohydrate Research, 145, 219-235 (1986)). Iota carrageenan, kappa carrageenan and lambda carrageenan are the most common forms but of these only iota and kappa carrageenan have gelling properties. Salts are

naturally present in the extracted carrageenans and commercial grades of carrageenan may contain added salts to increase gel strength.

5 WO 00/069275 in the name of Hercules Incorporated relates to the use of carrageenan as a gelatin replacement in sweetened food compositions.

WO 02/30214 in the name of C P Kelco US Inc relates to a gelatin-free gummy confection comprising gellan gum and nu-carrageenan, nu-/iota-carrageenan or both. No mention is made of liquid filled products.

10

EP-A-0 515 864 in the name of Hershey Foods Corporation relates to the use of a number of hydrocolloids as gelatin replacements whilst reducing the amount of fat in compositions such as caramel.

15 WO 02/30213 in the name of Société des Produits Nestlé SA relates to a food product comprising a gelatin-free water-based hydrocolloid casing enclosing a solid, liquid, soft or particulate core. The hydrocolloid may be carrageenan, alginate, agarose, gellan gum, pectin or a cellulose derivative.

20 It is an object of the present invention to provide a food product comprising a gelled water-based shell and a liquid centre wherein the shell avoids the use of gelatin and also has improved properties in terms of appearance, texture and ease of manufacture.

## SUMMARY OF THE INVENTION

25

The present invention provides a food product comprising a gelled water-based shell and a liquid centre characterised in that the water-based shell contains a mixture of kappa carrageenan and iota carrageenan in an amount sufficient to provide a gel texture, the relative proportions of kappa carrageenan and iota carrageenan being 50 to 30 95% by weight kappa carrageenan and 5 to 50% by weight iota carrageenan based on the sum of the weights of kappa and iota carrageenan.

The present invention also provides a process for the manufacture of a food product as defined above which comprises introducing the water-based shell composition in liquid form and the liquid centre into a mould so that the shell surrounds the centre and gelling the shell.

5

#### DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph showing the effect on gel temperature and viscosity of varying kappa carrageenan ratios in a shell composition.

Figure 2 is a graph showing optical density of the shell of liquid filled sweets.

10 Figure 3 is a graph of the influence of varying kappa carrageenan content on gel texture.

#### DETAILED DESCRIPTION OF THE INVENTION

15 The shell material is a water-based composition containing sufficient of a mixture of kappa carrageenan and iota carrageenan to provide a gel texture. The relative proportions of kappa carrageenan and iota carrageenan in this mixture should be 50 to 95% by weight kappa carrageenan and 5 to 50% by weight iota carrageenan based on the sum of the weights of kappa and iota carrageenan and such a mixture will be  
20 referred to hereinafter as "kappa/iota carrageenan".

The composition of the shell material is water-based and is prepared initially in liquid form. The amount of kappa/iota carrageenan present in the shell material is sufficient to gel the composition and provide the desired texture in the gelled material. The  
25 minimum amount of kappa/iota carrageenan can be determined experimentally in any particular case but has generally been found to be about 1.5% by weight. As the amount of kappa/iota carrageenan is increased, the gel becomes stiffer and it will not generally be appropriate to include more than 5% by weight. The amount of kappa/iota carrageenan in the shell will preferably be from 1.7 to 4% by weight, more  
30 preferably 2 to 2.5% by weight. It should be understood that the amount of kappa/iota carrageenan in the shell material is measured as pure kappa and iota carrageenan on a dry substance basis as a proportion of the shell solids. The relative proportions of kappa and iota carrageenan are calculated in the same way. Many commercially

available carrageenan preparations are not pure carrageenan but contain substantial amounts of other materials and this must be taken into account in any calculation.

5 Examples of the composition of suitable commercial sources of highly refined kappa and iota carrageenan are as follows:

kappa carrageenan: approximately 73% kappa and 4% iota

iota carrageenan: approximately 69% iota and 1% kappa.

10 The required blend of kappa/iota carrageenan can be prepared by mixing appropriate amounts of these materials. The proportions of kappa/iota carrageenan for use according to the present invention are as follows:

overall: 50 to 95% kappa, 5 to 50% iota

preferred: 60 to 95% kappa, 5 to 40% iota

more preferred: 65 to 90% kappa, 10 to 35% iota

most preferred: 70 to 85% kappa, 15 to 30% iota.

15 The composition to form the shell is formed as a blend in water and other main ingredients will be one or more sweeteners. The sweetener may be a sugar, a sugar syrup, corn syrup or a sugar substitute such as maltitol, lactitol, mannitol, xylitol, sorbitol, or an artificial sweetener. If desired the shell can be made sugar free by using  
20 only sugar substitute. The composition may also contain acid, acidity regulator and colourant as appropriate in the normal quantities. Water is added as needed to dissolve the ingredients before the shell material is cooked to the desired solids content for depositing. Further drying may take place after depositing. The final solids content of the shell may be from 50 to 90%, preferably from 60 to 85%, more preferably from 75  
25 to 85% by weight. The sweetener may be present in an amount of 45 to 88%, preferably 72 to 83% by weight based on the total weight of the final shell.

A water-based shell gelled with kappa/iota carrageenan will show a high level of transparency unless colourings or other materials are included which have the effect of  
30 reducing transparency. Preferably the shell is formulated in such a way that this natural transparency is retained. A 3.5 mm thick slice of the shell preferably has an optical density of 0.2 or less, more preferably 0.1 or less. A method for the measurement of the optical density of the shell is given below.

The liquid centre may be any appropriate composition for inclusion in a food product comprising a gelled water-based shell. The composition may be fat based such as a flavoured oil or chocolate or it may contain water and sweetener, for example a sugar, a sugar syrup, corn syrup or a sugar substitute such as maltitol, lactitol, mannitol, xylitol, sorbitol, or an artificial sweetener. If desired the liquid can be made sugar free by using only artificial sweetener and this may be particularly appropriate in combination with a sugar-free shell to provide an overall sugar free product. The liquid can contain other ingredients such as viscosifiers, humectants, colour, flavour, acid and/or functional ingredients such as minerals, vitamins and herbs. The water content of the composition will generally be from 10 to 30% by weight and the sweetener content from 70 to 90% by weight, in each case based on the total weight of the composition. The composition will generally have a viscosity in the range from 0.01 to 10,000 Pa s. The water activities of the shell and of the filling should be adjusted so that the filling remains liquid during the intended shelf life of the product.

The shell may have a thickness of 1mm to 10mm, preferably from 2mm to 7.5mm, more preferably from 2.5 to 5mm. The filling may comprise 5 to 90% of the overall product, preferably 7 to 30%, more preferably 10 to 25%.

The food product according to the invention may have any appropriate size or shape. Suitable shapes include spheres, hemispheres, cubes, teardrops, fruit shapes, animal shapes, insect shapes, pyramids and cylinders. The diameter (or largest dimension in the case of an irregularly shaped product) may be 10 to 50mm, preferably 12.5 to 40mm, more preferably 15 to 25mm.

The food product according to the present invention may be manufactured by a method which comprises introducing the water-based shell composition in liquid form and the liquid centre into a mould in such a manner that the shell surrounds the core and then gelling the shell. The shell composition may be introduced into the mould at elevated temperature and gelling takes place on cooling. For example the components may be introduced into starch moulds using a conventional filling apparatus such as a one shot Mogul head. The apparatus provides a co-axial deposit, initially depositing shell, then



introducing a flow of filling co-axially with the shell material and finishing with just shell. As an alternative to starch moulds, metal or plastic moulds could also be used.

5 The use of kappa/iota carrageenan has surprisingly been found to produce good shell properties combined with ease of processing when used with the liquid filled food products according to the invention.

10 Kappa carrageenan is generally thought to produce a brittle gel so that using 50% or more kappa carrageenan would not have been expected to provide a gel with an acceptable texture. However, it has surprisingly been found that when a gel made using kappa/iota carrageenan is used as the shell for a liquid filled product such as a sweet, the texture is perceived as pleasant and the brittleness is even an advantage as it enhances the contrast in mouth-feel between the shell and the liquid. The whole sweet is perceived by the consumer as being chewy with bouncy characteristics.

15 As already noted, a shell formed from a composition gelled with kappa/iota carrageenan is transparent in the absence of ingredients that reduce its natural transparency and this is in contrast to gelatin/starch gels which tend to be opaque as a result of the presence of the starch in formulations which give suitable eating texture and processability. The transparency of the shell is a particular advantage in terms of  
20 the appearance of the product since it enables the consumer to see the filling which adds to the appeal of the product. A clearly visible filling is particularly advantageous when the filling contains active ingredients such as vitamins, minerals or herbs.

25 It is a particular advantage of the use of kappa/iota carrageenan that the advantageous properties can be achieved without the use of gelatin. Although the presence of gelatin is not absolutely excluded there is no particular reason why it is required and, in view of the disadvantages associated with the use of gelatin referred to above, it is preferred that the shell is gelatin free.

30

The use of kappa/iota carrageenan provides particular advantages in terms of viscosity in that the relatively high content of kappa carrageenan produces a liquid of relatively low viscosity. This is a particular advantage in terms of uniform free flow of the

product into the mould and producing a uniform product with a well centred filling. In addition, tailing (the tendency for a trail of shell material to be deposited as the depositing head moves from one mould to another) is easier to avoid when using a shell composition of lower viscosity which makes high speed production of the products easier to achieve.

A further advantage of kappa/iota carrageenan is that the core material sets relatively quickly and, in particular, much more quickly than starch/gelatin based gels. Whilst the shell remains liquid, there is a tendency for the liquid filling to migrate to the edge of the shell which results in the filling being poorly centered. The fact that the shell is thin in places means that the filling has a tendency to leak during handling and storage of the product. Better centering of the filling and a reduced tendency to leak means that the shell can be made thinner and the filling can make up a larger proportion of the overall product. This further enhances the visibility of the filling, with the advantages discussed above.

## EXAMPLES

The invention is illustrated by the following examples.

### Example 1

Shell compositions were made to the following recipe, the amounts being in grams:

	water	1800
25	citric/malic acids (powder)	30
	acidity regulator (sodium citrate)	10
	high fructose corn syrup	800
	42DE glucose syrup	2100
	carrageenan	100
30	sucrose	2200

Three different compositions were made with the following carrageenens:

~~52% commercial kappa-carrageenan 48% commercial iota-carrageenan (52%  
kappa/48% iota based on the total of pure kappa and iota)~~

75% commercial kappa-carrageenan 25% commercial iota-carrageenan (73%  
kappa/27% iota based on the total of pure kappa and iota)

5 100% commercial kappa-carrageenan (95% kappa/5% iota based on the total of pure  
kappa and iota).

The three compositions referred to above were cooked to a total solids content of 77%  
and then deposited into starch moulds at 90°C using a centre-in-shell depositor with a  
10 centre having the following composition:

sugar syrup	99.5%
colour	0.004%
flavour	0.4%.

The starch impressions were in the shapes of insects and animals as for the Rowntrees  
15 Bursting Bugs product. The filling was 12% by weight of the finished sweets. The  
sweets were cooled to 20°C in their starch trays and allowed to dry for 24 hours before  
being demoulded. The final total solids content of the shell was 79%.

Eating the sample with 52% commercial kappa-carrageenan and 48% commercial iota-  
20 carrageenan did not give a particularly strong liquid centre impression. The shell had a  
soft texture and so, when it was broken by the teeth, there was only a small contrast  
between the liquid filling and the pieces of shell.

With the sample 75% commercial kappa-carrageenan and 25% commercial iota-  
25 carrageenan, the moment at which the shell broke to release the filling was more clearly  
felt. This was due to the textural contrast between the shell and the liquid, making the  
liquid more noticeable in the mouth. Once the liquid had cleared from the mouth, the  
pieces of shell which remained were sufficiently cohesive to still be chewy and pleasant  
to eat.

30

The liquid release could clearly be felt in sample 100% commercial kappa-carrageenan,  
but once the liquid had cleared from the mouth, the fragments of shell remaining were  
brittle and gave a less pleasant eat.

### Example 2

Viscosity as a function of temperature was measured for the three shell compositions referred to in Example 1 using a Haake RS150 rheometer with concentric cylinder measuring system. The shear rate was constant (set at  $1\text{ s}^{-1}$ ) and the temperature ramp was  $1^\circ\text{C}/\text{minute}$ . The results are shown in Figure 1. Viscosity decreases as % kappa increases and the gel temperature (inflection point indicated with an arrow) reduces slightly with increasing kappa.

### Example 3

- Optical density of the shell of the product from example 1 (75% commercial kappa-carrageenan, 25% commercial iota-carrageenan) was measured and compared to the optical densities of the shells of the following other confectionery sweets:

Rowntrees Red Bursting Bug

Rowntrees Orange Bursting Bug

- Rowntrees Green Bursting Bug

Betty Crocker Fruit Gushers Fruitomic Punch Dark Blue

Betty Crocker Fruit Gushers Fruitomic Punch Green

Betty Crocker Fruit Gushers Fruitomic Punch Yellow

- The samples of the other sweets were purchased through normal commercial channels in March 2003.

The method was as follows:

- A special cutting device was made by gluing two razor blades together with spacers placed in between the blades so that the blades were a fixed distance apart and the cutting edges parallel. Using callipers the blades were found to be  $3.50 \pm 0.05\text{ mm}$  apart. These blades were then used to cut slices having a uniform thickness from the shell of the liquid filled confectionery sweets. The slice was then sandwiched in between two glass microscope slides.

30

The optical density of each gel slice was measured using a GRETAG MACBETH D200-11 Transmission Densitometer. Before any measurements were taken the machine was calibrated using the method described in the operating instructions. The

3mm measuring aperture was placed in the machine, and the machine set to the Density Measurement mode (Mode '1'). The zero level was then set with two microscope slides in the optical path. The two slides used for zero calibration were then replaced by the gel slice sample held in between two glass slides. The optical density of the sample was then measured in the manner described in the instruction manual.

Four measurements were taken of each slice in different positions avoiding the edges, and slices from two different sweets were measured. The results were averaged and plotted together with error bars of 2 x standard deviation. The results are shown in Figure 2.

#### Example 4


Confectionery sweets were prepared according to the shell recipe of Example 1 but using carrageenan having the following proportions of kappa to iota:

- 100% commercial kappa-carrageenan
- 75% commercial kappa-carrageenan 25% commercial iota-carrageenan
- 60% commercial kappa-carrageenan 40% commercial iota-carrageenan
- 50% commercial kappa-carrageenan 50% commercial iota-carrageenan.

The sweets were moulded into cylindrical starch impressions with no filling and the sweets were 8mm tall with a diameter of 17.5mm.

The texture of the sweets was measured using a Stable Micro Systems TA-HD texture analyser fitted with a 40mm diameter cylindrical probe. Two compressions were made to a compression of 60% strain with a 5s time interval between the compressions. The probe speed was 1mm/s. 6 examples of each carrageenan ratio were analysed and values for hardness, cohesiveness and resilience were recorded.

The calculation of hardness, cohesiveness and resilience is shown in the following figure:



Hardness : Force F1

20 Cohesiveness:  $\text{Area B} / (\text{Area A1} + \text{Area A2})$

Resilience:  $\text{Area A2} / \text{Area A1}$

Hardness is the force required to reach the defined percentage of strain. It is measured as the peak force during the first compression.

25

Cohesiveness is related to the work required to overcome the internal structure of the sample. It is calculated as the ratio of the positive area during the second compression to that during the first compression.

30 Resilience or elasticity corresponds to a measure of how rubber-like the sample is. It is calculated as the ratio of the area during recovery to the initial state from the first peak force to that required to reach the first peak force.

35 The results are shown in Figure 3 and they show that the texture of the shell changes as the amount of kappa carrageenan increases and, in particular, it becomes harder but resilience and cohesiveness decrease. This texture change leads to a shell which enhances the impact of the liquid filling in the mouth, the shell breaks easily when

chewed to release the liquid and the hard texture of the shell has good contrast with the liquid filling when mixed in the mouth.

#### Composition of Carrageenens

- 5 As noted above, the carrageenens used in the above examples were commercial highly refined kappa-carrageenan and commercial highly refined iota-carrageenan.

Moisture contents were measured by gravimetric determination of mass loss during oven drying at 102°C for 4 hours.

Protein was determined using the Kjeldahl method with a conversion factor of 6.25.

- 10 Cation (Ca, Mg, Na, K) and sulphur determination was performed by ICP-AES (optical emission spectrometry with inductively coupled plasma) after microwave digestion. Assuming that the sulphur originates exclusively from carrageenan-bound sulphate, the total sulphate content was then recalculated.

Free sugars were determined by HPLC (high performance liquid chromatography).

- 15 The composition of these carrageenans was determined by  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR at the NIZO Food Research Institute (Netherlands). The sample preparation and the analysis conditions are described below.

#### $^1\text{H}$ -NMR

20

The samples were dissolved at 70°C in 20mM  $\text{Na}_2\text{HPO}_4/\text{D}_2\text{O}$  to a concentration of about 0.8 % (w/v) and sonicated. The NMR spectra were recorded at 70°C on a Varian INOVA 600 spectrometer operating at 599.93 MHz. The interpretation of the spectra is based on the resonances of the anomeric protons of carrageenans between 5 and 5.7

- 25 ppm. The respective chemical shifts of kappa-, iota-, mu-, nu- and lambda-carrageenans are known, the other peaks in this area are designated as unidentified.

The unidentified material may be desulphated carrageenans, or even not a carrageenan material at all. It will not, however, be kappa-, iota-, mu-, nu- or lambda-carrageenan as these are determined by their known peaks. The carrageenan content is calculated as

- 30 a molar percentage of the total resonances in this spectral region (excluding glucose and sucrose). Citrate can be identified by  $^1\text{H}$ -NMR but was not detected.

<sup>13</sup>C-NMR

The samples were dissolved in H<sub>2</sub>O, sonicated, dialysed against 10 mM NaHPO<sub>4</sub> and distilled water. Then, they were lyophilised and dissolved in D<sub>2</sub>O to a concentration of about 8% (w/v). The NMR spectra were recorded at 74°C on a Bruker AMX500 spectrometer operating at 125.76 MHz. Typically 40000 scans were accumulated using an inter-pulse delay of 1.5s and WALTZ-16 proton decoupling. <sup>1</sup>H-NMR spectra taken before and after the <sup>13</sup>C-NMR experiment were identical for both samples, indicating that the samples were stable during the <sup>13</sup>C-NMR experiment.

## Calculation

The total carrageenan by product weight was estimated assuming that "non-carrageenan" and "total carrageenan" represent 100% of the product content. The "non-carrageenan" was obtained by summing up the moisture, proteins, free sugars, K, Na, Mg, and Ca contents. The weight by product weight content of the different types of carrageenans was obtained using the mole % values and recalculating the proportion of each type relative to the total carrageenans.

The results were as follows:

Commercial kappa-carrageenan

## Carrageenans

kappa (% w/w)	73
iota (% w/w)	4
mu (% w/w)	2
nu (% w/w)	n.d.
unidentified (% w/w)	2

The remainder was moisture (around 12%), protein and salts. No free sugars were detected.



Commercial iota-carrageenan

## Carrageenans

	kappa (% w/w)	1
5	iota (% w/w)	69
	mu (% w/w)	n.d.
	nu (% w/w)	n.d.
	unidentified (% w/w)	9

- 10 The remainder was moisture (around 12%), protein and salts. No free sugars were detected.

(n.d. = not detected)

## CLAIMS:

1. A food product comprising a gelled water-based shell and a liquid centre characterised in that the water-based shell contains a mixture of kappa carrageenan and  
5 iota carrageenan in an amount sufficient to provide a gel texture, the relative proportions of kappa carrageenan and iota carrageenan being 50 to 95% by weight kappa carrageenan and 5 to 50% by weight iota carrageenan based on the sum of the weights of kappa and iota carrageenan.
- 10 2. A food product as claimed in claim 1 wherein the shell contains, on a dry substance basis, 1.5% to 5% by weight of the mixture of kappa and iota carrageenans.
3. A food product as claimed in claim 1 wherein the shell contains, on a dry substance basis, 1.7% to 4% by weight of the mixture of kappa and iota carrageenans.
- 15 4. A food product as claimed in claim 1 wherein the shell contains, on a dry substance basis, 2% to 2.5% by weight of the mixture of kappa and iota carrageenans.
5. A food product as claimed in any of claims 1 to 4 wherein the mixture of kappa  
20 and iota carrageenans contains 60 to 95% kappa and 5 to 40% iota based on the total weight of kappa and iota.
6. A food product as claimed in any of claims 1 to 4 wherein the mixture of kappa  
25 and iota carrageenans contains 65 to 90% kappa and 10 to 35% iota based on the total weight of kappa and iota.
7. A food product as claimed in any of claims 1 to 4 wherein the mixture of kappa  
and iota carrageenans contains 70 to 85% kappa and 15 to 30% iota based on the total weight of kappa and iota.
- 30 8. A food product as claimed in any of claims 1 to 7 wherein the shell is formed of a composition comprising in addition to the blend of kappa and iota carrageenan, water and one or more sweeteners.

9. A food product as claimed in claim 8 wherein the sweetener is a sugar, a sugar syrup, corn syrup or a sugar substitute or an artificial sweetener.

5 10. A food product as claimed in claim 8 or 9 wherein the sweetener is present in an amount of 45 to 88% of the shell.

11. A food product as claimed in claim 8 or 9 wherein the sweetener is present in an amount of 72 to 83% of the shell.

10

12. A food product as claimed in any of claims 1 to 11 wherein the composition forming the shell also contains acid, acidity regulator and/or colourant.

13. A food product as claimed in any of claims 1 to 12 wherein the solids content of  
15 the shell is 50 to 90% by weight.

14. A food product as claimed in any of claims 1 to 12 wherein the solids content of the shell is 60 to 85% by weight.

20 15. A food product as claimed in any of claims 1 to 12 wherein the solids content of the shell is 75 to 85% by weight.

16. A food product as claimed in any of claims 1 to 15 wherein a 3.5mm thick slice of the shell has an optical density of 0.2 or less.

25

17. A food product as claimed in any of claims 1 to 15 wherein a 3.5mm thick slice of the shell has an optical density of 0.1 or less.

18. A food product as claimed in any of claims 1 to 17 wherein the liquid centre is a  
30 fat based composition.

19. A food product as claimed in claim 18 wherein the liquid centre is a flavoured oil or chocolate.

20. A food product as claimed in any of claims 1 to 19 wherein the liquid centre contains water and sweetener.

5 21. A food product as claimed in claim 20 wherein the sweetener is a sugar, a sugar syrup, corn syrup or a sugar substitute or an artificial sweetener.

22. A food product as claimed in claim 21 wherein the liquid centre contains colour, flavour, acid and/or one or more functional ingredients.

10

23. A food product as claimed in claim 22 wherein the liquid centre contains at least one of minerals, vitamins and herbs.

15

24. A food product as claimed in any of claims 20 to 23 wherein the water content of the liquid centre is from 10 to 30% by weight and the sweetener content is from 70 to 90% by weight.

20

25. A food product as claimed in any of claims 1 to 24 wherein the viscosity of the liquid centre is in the range from 0.01 to 10,000 Pa s.

26. A food product as claimed in any of claims 1 to 25 wherein the shell has a thickness of 1mm to 10mm.

25

27. A food product as claimed in any of claims 1 to 25 wherein the shell has a thickness of 2mm to 7.5mm.

28. A food product as claimed in any of claims 1 to 25 wherein the shell has a thickness of 2.5 to 5mm.

30

29. A food product as claimed in any of claims 1 to 28 wherein the liquid filling comprises 5 to 90% of the product.

30. A food product as claimed in any of claims 1 to 28 wherein the liquid filling comprises 7 to 30% of the product.
31. A food product as claimed in any of claims 1 to 28 wherein the liquid filling comprises 10 to 25% of the product.
32. A food product as claimed in any of claims 1 to 31 which is gelatin free.
33. A process for the manufacture of a food product as claimed in any of claims 1 to 32, which comprises introducing the water-based shell composition in liquid form and the liquid centre into a mould so that the shell surrounds the centre and gelling the shell.
34. A process as claimed in claim 33 wherein the components are introduced into starch moulds.
35. A process as claimed in claim 33 or 34 wherein the water-based shell composition in liquid form and the liquid centre are introduced into a mould using an apparatus which provides a co-axial deposit, initially depositing shell, then introducing a flow of filling co-axially with the shell material and finishing with just shell.

## ABSTRACT

## GELLED FOOD PRODUCT WITH A LIQUID CENTRE

- 5 The invention provides a food product comprising a gelled water-based shell and a liquid centre characterised in that the water-based shell contains a mixture of kappa carrageenan and iota carrageenan in an amount sufficient to provide a gel texture, the relative proportions of kappa carrageenan and iota carrageenan being 50 to 95% by weight kappa carrageenan and 5 to 50% by weight iota carrageenan based on the sum  
10 of the weights of kappa and iota carrageenan. The shell avoids the use of gelatin and has improved properties in terms of appearance, texture and ease of manufacture.

Figure 1

Effect on gel temperature and viscosity of varying Kappa Carrageenan ratios  
shear rate 1s-1




Figure 2

Optical Density of Liquid Filled Gummie Shells






Figure 3





**PCT/EP2004/002490**



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**